## Pyruvic acid-<sup>13</sup>C sodium

Cat. No.:	HY-Y0781S	•
CAS No.:	87976-70-3	Q
Molecular Formula:	C <sub>2</sub> <sup>13</sup> CH <sub>3</sub> NaO <sub>3</sub>	
Molecular Weight:	111.04	130
Target:	Endogenous Metabolite	'°Ç ONa
Pathway:	Metabolic Enzyme/Protease	
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)	0

BIOLOGICAL ACTIVITY		
Description	Pyruvic acid- <sup>13</sup> C (sodium) is the <sup>13</sup> C-labeled Pyruvic acid. Pyruvic acid is an intermediate metabolite in the metabolism of carbohydrates, proteins, and fats.	
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

## REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

[2]. Kładna A, et al. Reactivity of pyruvic acid and its derivatives towards reactive oxygen species. Luminescence. 2015 Nov;30(7):1153-8.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

**Product** Data Sheet

