Inhibitors

MCE RedChemExpress

Levulinic acid-d₅

Cat. No.: HY-Y0839S CAS No.: 1206185-52-5 Molecular Formula: $C_5H_3D_5O_3$ Molecular Weight: 121.15

Target: Endogenous Metabolite

Pathway: Metabolic Enzyme/Protease

Storage: Pure form -20°C 3 years

 $\begin{array}{ccc} & 4^{\circ}\text{C} & 2 \text{ years} \\ \text{In solvent} & -80^{\circ}\text{C} & 6 \text{ months} \\ & -20^{\circ}\text{C} & 1 \text{ month} \end{array}$

BIOLOGICAL ACTIVITY

Description	Levulinic acid- d_5 is the deuterium labeled Levulinic acid[1]. Levulinic acid is a precursor for the synthesis of biofuels, such as ethyl levulinate[2].
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

 $[2]. \ Pileidis\ FD, et\ al.\ Levulinic\ Acid\ Biorefineries:\ New\ Challenges\ for\ Efficient\ Utilization\ of\ Biomass.\ ChemSus\ Chem.\ 2016\ Mar\ 21;9(6):562-82.$

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA