


## Glycine

<b>Cat. No.:</b>	HY-Y0966		
<b>CAS No.:</b>	56-40-6		
<b>Molecular Formula:</b>	C <sub>2</sub> H <sub>5</sub> NO <sub>2</sub>		
<b>Molecular Weight:</b>	75.07		
<b>Target:</b>	Endogenous Metabolite; iGluR		
<b>Pathway:</b>	Metabolic Enzyme/Protease; Membrane Transporter/Ion Channel; Neuronal Signaling		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

H<sub>2</sub>O : 6 mg/mL (79.93 mM; Need ultrasonic)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	13.3209 mL	66.6045 mL	133.2090 mL
5 mM	2.6642 mL	13.3209 mL	26.6418 mL
10 mM	1.3321 mL	6.6605 mL	13.3209 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

Glycine is an inhibitory neurotransmitter in the CNS and also acts as a co-agonist along with glutamate, facilitating an excitatory potential at the glutamergic N-methyl-D-aspartic acid (NMDA) receptors.

#### IC<sub>50</sub> & Target

Human Endogenous Metabolite

### REFERENCES

[1]. Johnson JW, et al. Glycine potentiates the NMDA response in cultured mouse brain neurons. Nature. 1987 Feb 5-11;325(6104):529-31.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA