


Fmoc-Gly-OH-2-¹³C,¹⁵N

| | | | |
|---------------------------|--|-------|----------|
| Cat. No.: | HY-Y1250S1 | | |
| CAS No.: | 285978-12-3 | | |
| Molecular Formula: | C ₁₆ ¹³ CH ₁₅ ¹⁵ NO ₄ | | |
| Molecular Weight: | 299.29 | | |
| Target: | Isotope-Labeled Compounds | | |
| Pathway: | Others | | |
| Storage: | Powder | -20°C | 3 years |
| | | 4°C | 2 years |
| | In solvent | -80°C | 6 months |
| | | -20°C | 1 month |



SOLVENT & SOLUBILITY

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (6.95 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: 2.08 mg/mL (6.95 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.08 mg/mL (6.95 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Fmoc-Gly-OH-2-¹³C,¹⁵N is the ¹³C,¹⁵N labeled Fmoc-Gly-OH-2[1].

In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA