

MyD88

MyD88 (Myeloid differentiation primary response gene 88) is a protein that, in humans, is encoded by the MYD88 gene. Available evidence suggests that MYD88 is dispensable for human resistance to common viral infections and to all but a few pyogenic bacterial infections, demonstrating a major difference between mouse and human immune responses. MyD88 is an essential adaptor protein in the IL-1R1 signaling pathway. MyD88 may define a family of signal transduction molecules with an ancestral function in the activation of the immune system. MyD88 functions as a pure adaptor linking the IL-1R1 to downstream IRAK kinases. Mutation in MYD88 at position 265 leading to a change from leucine to proline have been identified in many human lymphomas including ABC subtype of Diffuse Large B-cell Lymphoma and Waldenstrom's Macroglobulinemia.

MyD88 Inhibitors

Schaftoside

Cat. No.: HY-N0703

Schaftoside is a flavonoid found in a variety of Chinese herbal medicines, such as Eleusine indica. Schaftoside inhibits the expression of TLR4 and Myd88. Schaftoside also decreases Drp1 expression and phosphorylation, and reduces mitochondrial fission.

Purity: 99.88%

Clinical Data: No Development Reported Size: 10 mM × 1 mL, 5 mg, 10 mg, 20 mg

T6167923

Cat. No.: HY-19744

T6167923 is a potent and selective inhibitor of MyD88-dependent signaling pathways. T6167923 directly binds to Toll/IL1 receptor (TIR) domain of MyD88 and disrupts MyD88 homodimeric formation.

Purity: 99.08%

Clinical Data: No Development Reported

10 mM × 1 mL, 5 mg, 10 mg, 25 mg, 50 mg, 100 mg

TLR1

Cat. No.: HY-W011400

TLR1 (compound 4a) is a low molecular weight, cell-penetrating Toll/IL-1 receptor/resistance (TIR) domain/BB-Loop mimic. TLR1 inhibits IL-1 receptor-mediated responses.

Purity: ≥99.0%


Clinical Data: No Development Reported Size: 500 μg (33 mM * 50 μL in Ethanol)

ST 2825

ST 2825 is a specific MyD88 dimerization inhibitor, ST2825 interferes with recruitment of IRAK1 and IRAK4 by MyD88, causing inhibition of IL-1β-mediated activation of NF- κ B transcriptional activity.

Purity: 99.86%

Clinical Data: No Development Reported Size: 10 mM × 1 mL, 1 mg, 5 mg, 10 mg



TJ-M2010-5

TJ-M2010-5 is a MyD88 inhibitor that binds to the TIR domain of MyD88 to interfere with its homodimerization, and the TLR/MyD88 signal pathway. TJ-M2010-5 can be used for the research of myocardial ischemia/reperfusion injury (MIRI).

Cat. No.: HY-139397

99.25%

Clinical Data: No Development Reported

5 mg, 10 mg, 25 mg, 50 mg, 100 mg