

100 bp DNA Marker


1 Contents

Components	HY-K0802-250 μ L	HY-K0802-500 μ L	HY-K0802-1 mL
100 bp DNA Marker	250 μ L	250 μ L \times 2	250 μ L \times 4

2 Introduction

The 100 bp DNA Marker is provided in a solution of 1 \times DNA Loading Buffer, which can be directly used for nucleic acid electrophoresis analysis. The Marker contains 10 double-stranded DNA fragments ranging from 100 bp to 1000 bp. 5 μ L of this product contains about 120 ng for the 500 bp band, and about 50 ng for the other bands.

3 Electrophoresis illustration



1.5% Agarose 0.5x TBE Buffer
5 μ L/lane 7 V/cm, 40 min

4 Protocol

1. Add 5 μ L of 100 bp DNA Marker to sample well of the agarose gel and perform electrophoresis.
2. After electrophoresis, stain with Nucleic Acid Gel Stain and detect the electrophoresis results.

Note: a) 1.2-2.0% agarose gel at 5-10 V/cm is recommended.

- b) Adjust the loading volume of DNA Marker for different loading well format.
- c) Pre-dyeing or post-dyeing is suitable when using the Nucleic Acid Gel Stain.

5 Storage

-20°C, 2 years.

Avoid repetitive freeze-thaw cycles.

6 Precautions

1. For short-term use, DNA Marker may be stored at 2-8°C.
2. Replace the electrophoresis buffer in time and use fresh agarose gels to achieve better electrophoretic results.
3. When the concentration of agarose gel is too low, the bands will not be easy to separate.
4. This product is for R&D use only, not for drug, household, or other uses.
5. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

7 Recommended products used for Nucleic Acid Gel Electrophoresis

Cat. No	Name	Application
HY-K1031	Agarose	Agarose gel
HY-K1029	Agarose With TAE Powder (1%)	
HY-K1016	TBE Powder (1 L of 1x)	
HY-K1015	TAE Powder (1 L of 1x)	Electrophoresis buffer
HY-K1017	Rapid Running Buffer Powder (1 L of 1x)	
HY-K1004	SYBR Green I Nucleic Acid Gel Stain	Nucleic Acid Gel Stain
HY-K1007	Red Nucleic Acid Gel Stain (10,000x)	