Tel: 609-228-6898

Email: tech@MedChemExpress.com



15 kb DNA Marker


1 Contents

| Components | HY-K0808-250 μL | HY-K0808-500 μL | HY-K0808-1 mL |
|------------------|-----------------|-----------------|---------------|
| 15 kb DNA Marker | 250 μL | 250 μL × 2 | 250 μL × 4 |

2 Introduction

The 15 kb DNA Marker is provided in a solution of $1 \times$ DNA Loading Buffer, which can be directly used for nucleic acid electrophoresis analysis. The Marker contains 8 double-stranded DNA fragments ranging from 500 bp to 15 kb. 5 μ L of this product contains about 80 ng for the 4000 bp band, and about 30 ng for the other bands.

3 Electrophoresis illustration



0.7% Agarose 1x TAE Buffer $5 \mu L/lane 7 V/cm, 40min$

4 Protocol

- 1. Add 5 μ L of 15 kb DNA Marker to sample well of the agarose gel and perform electrophoresis.
- 2. After electrophoresis, stain with Nucleic Acid Gel Stain and detect the electrophoresis results.

Note: a) 0.7-1.0% agarose gel at 5-10 V/cm and 1× TAE Powder are recommended.

- b) Adjust the loading volume of DNA Marker for different loading well format.
- c) Pre-dyeing or post-dyeing is suitable when using the Nucleic Acid Gel Stain.

Storage

-20°C, 2 years.

Avoid repetitive freeze-thaw cycles.

Precautions

- 1. For short-term use, DNA Marker may be stored at 2-8°C.
- 2. Replace the electrophoresis buffer in time and use fresh agarose gels to achieve better electrophoretic results.
- 3. When the concentration of agarose gel is too high (no more than 1.0 %), the bands will not be easy to separate. TBE Powder is not recommended.
- 4. This product is for R&D use only, not for drug, household, or other uses.
- 5. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Recommended products used for Nucleic Acid Gel Electrophoresis

| Cat. No | Name | Application | |
|----------|---|------------------------|--|
| HY-K1031 | Agarose | Agarose gel | |
| HY-K1029 | Agarose With TAE Powder (1%) | | |
| HY-K1016 | TBE Powder (1 L of 1×) | | |
| HY-K1015 | TAE Powder (1 L of 1×) | Electrophoresis buffer | |
| HY-K1017 | Rapid Running Buffer Powder (1 L of 1×) | | |
| HY-K1004 | SYBR Green I Nucleic Acid Gel Stain | Nucleic Acid Gel Stain | |
| HY-K1007 | Red Nucleic Acid Gel Stain (10,000x) | | |

Tel: 609-228-6898