

Sbf I

1 Contents

| Component | HY-KE7036-25T |
|------------------|---------------|
| Sbf I | 25 µL |
| 10× Buffer | 1 mL |
| 10× Color Buffer | 1 mL |

2 Introduction

Sbf I is a restriction enzyme for rapid DNA digestion, including plasmid, genomic DNA as well as PCR products. Isoschizomers: Sda I, Sse8387 I. Sbf I is 100% active in 10× Buffer and 10× Color Buffer. 10× Color Buffer includes a density reagent along with red and yellow tracking dyes that allow for direct loading of the reaction mixtures on a gel. The red dye of the Color Buffer migrates with 2500 bp DNA fragments in a 1% agarose gel and the yellow dye of the Color Buffer migrates faster than 10 bp DNA fragments in a 1% agarose gel.

Cleavage site



3 General Protocol

1. Fast Digestion of Different DNA

1.1 Combine the following reaction components on ice in the order indicated:

| | Plasmid DNA | PCR product | Genomic DNA |
|--------------------------------|-------------------|------------------|--------------|
| ddH ₂ O | 15 µL | 16 µL | 30 µL |
| 10× Buffer or 10× Color Buffer | 2 µL | 3 µL | 5 µL |
| DNA | 2 µL (up to 1 µg) | 10 µL (~ 0.2 µg) | 10 µL (5 µg) |
| Sbf I | 1 µL | 1 µL | 5 µL |
| Total | 20 µL | 30 µL | 50 µL |

Note: When PCR product will be used for cloning, it is recommended to purify PCR product prior digestion.

1.2 Mix gently and spin down.

1.3 Incubate at 37°C for 15 min (Plasmid DNA), or for 15-30 min (PCR product), or for 30-60 min (Genomic DNA).

1.4 (Optional) Inactivate the enzyme by heating for 20 min at 80°C.

1.5 If 10× Color Buffer was used in the reaction, load an aliquot of the reaction mixture directly on a gel.

2. Scaling up Plasmid DNA Digestion Reaction

| | | | | | |
|--------------------------------|-------|-------|-------|-------|-------|
| DNA | 1 µg | 2 µg | 3 µg | 4 µg | 5 µg |
| Sbf I | 1 µL | 2 µL | 3 µL | 4 µL | 5 µL |
| 10× Buffer or 10× Color Buffer | 2 µL | 2 µL | 3 µL | 4 µL | 5 µL |
| Total | 20 µL | 20 µL | 30 µL | 40 µL | 50 µL |

Note: Increase the incubation time by 3-5 min if the total reaction volume exceeds 20 µL. It is recommended to use heat block or water thermostat.

4 Number of Recognition Sites in DNA

| λDNA | ΦX174 | pBR322 | pUC57 | pUC18/19 | SV40 | M13mp18/19 | Adeno2 |
|------|-------|--------|-------|----------|------|------------|--------|
| 5 | 0 | 0 | 0 | 1 | 0 | 1 | 3 |

5 Methylation Effects on Digestion

| Dam | Dcm | CpG | EcoKI | EcoBI |
|-----------|-----------|-----------|-----------|-----------|
| no effect | no effect | no effect | no effect | no effect |

6 Storage

-20°C, 2 years.

7 Precautions

1. The combined volume of the enzymes in the reaction mixture should not exceed 1/10 of the total reaction volume.
2. Use 1 µL of each enzyme and scale up the reaction conditions appropriately.
3. If the enzymes require different reaction temperatures, start with the enzyme that requires a lower temperature, then add the second enzyme and incubate at the higher temperature.
4. This product is for R&D use only, not for drug, household, or other uses.
5. For your safety and health, please wear a lab coat and disposable gloves to operate.